

PLANT POLYSACCHARIDE DEGRADATION BY FUNGI

RONALD P. DE VRIES¹, AD WIEBENGA¹, PEDRO M. COUTINHO², BERNARD HENRISSAT²

¹CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands

²AFMB, 10UMR 6098 CNRS-Universités Aix-Marseille I& II, 13288 Marseille Cedex 9, France

r.devries@cbs.knaw.nl

ABSTRACT

Plant biomass is the major carbon source for many fungal species. Due to its complex polymeric nature, degradation of this biomass to digestible monomers requires a large range of enzyme activities. With the availability of an increasing number of fungal genomes, new insights into the diversity of fungi with respect to plant biomass degradation have been obtained. Recent progress in this area will be reviewed in this paper.

Keywords: fungi, plant biomass, polysaccharide degradation, CAZy, growth profiling

PLANT BIOMASS COMPOSITION AND STRUCTURE

Plant biomass is the most abundant organic matter on earth and the major substrate for the majority of fungal species. It consists mainly of polysaccharides, but also contains proteins and the aromatic polymer lignin. Plant polysaccharides can be divided into plant cell wall polysaccharides (cellulose, hemicelluloses, pectin) and storage polysaccharides (e.g. starch, inulin, gums) [1]. They consist of many different monomeric components that are attached to each other by a variety of linkages.

Cellulose is the most abundant plant polysaccharide. It consists of a linear chain of β -1,4-linked D-glucose residues (Fig. 1). These chains are organised in bundles called microfibrils [2] that provide the main strength and structure for the plant cell wall.




Figure 1: Schematic presentation of cellulose, its degradation and release of D-glucose. Reprinted from [3] with permission from the publisher (Copyright remains with the original publisher).

Hemicelluloses are more diverse in nature and three main types are present in plant cell walls: xylan, xyloglucan and mannan (Fig. 2). Xylan consists of a backbone of β -1,4-linked D-xylose residues [4, 5]. Other residues can be attached to this backbone, such as α -linked L-arabinose or (4-O-methyl-) D-glucuronic acid residues, α - or β -linked D-galactose residues and acetyl residues [5-8]. The L-arabinose residues can be further decorated with feruloyl residues [9-11]. Xyloglucan has a backbone that consists of β -1,4-linked D-glucose residues that are decorated with α -1,6-linked D-xylose residues [12]. Attached to the D-xylose residues can be L-arabinose, D-galactose and L-fucose residues. Mannan is often also referred to as galactomannan as this

hemicelluloses consists of a of β -1,4-linked D-mannose backbone with α -1,4-linked D-galactose residues and acetyl residues as decorations [13]. Depending on the plant species, the D-mannose backbone can also be interrupted with single D-glucose residues.




Figure 2: Schematic presentation of hemicellulose components (xylan, galacto(gluco)mannan, xyloglucan) and their degradation. Reprinted from [3] with permission from the publisher (Copyright remains with the original publisher).

Pectin is the third plant cell wall polysaccharide and it consists of several sub-structures (Fig. 3) [14, 15]. Homogalacturonan is a linear polymer of α -1,4-linked D-galacturonic acid residues that has various degrees of acetylation and/or methylation. Xylogalacturonan is a modified homogalacturonan with β -1,3-linked D-xylose residues attached to D-galacturonic acid. Rhamnogalacturonan 1 consists of a backbone of α -1,4-linked D-galacturonic acid and α -1,2-linked L-rhamnose residues. Side chains consisting of L-arabinose and/or D-galactose residues are attached to many of the L-rhamnose residues creating a structure often referred to as the 'hairy' region of pectin. Terminal feruloyl and *p*-coumaroyl residues can be attached to these side chains.

Starch is one of the main storage polysaccharides. It consists of an α -1,4-linked polymer of D-glucose residues that contain α -1,4-linked branching points [16]. Another major storage polysaccharide is inulin, that consist of a β -2,1-linked chain of D-fructose with a terminal d-glucose residue [17]. Another varied group of storage polysaccharides are the gums that contain many different structures [18]. Some gums (e.g. locust bean gum, guar gum) are highly similar in structure to cell wall galactomannan.




Figure 3: Schematic presentation of pectin components and their degradation. Reprinted from [3] with permission from the publisher (Copyright remains with the original publisher).

BIODEGRADATION OF PLANT POLYSACCHARIDES

Due to the diverse nature of plant polysaccharides, a large range of enzyme activities is required to degrade them to their monomeric components (Table 1) [1, 19]. These enzymes can be divided into families based on modules in their amino acid sequence resulting in a classification system called Carbohydrate Active enZymes (CAZy, www.cazy.org) [20]. Using the CAZy annotation pipeline for fungal genomes has been shown to be a highly powerful tool to obtain insight in the carbohydrate potential of a fungus and has been shown to be superior to manually annotated carbohydrate related genes of fungal genomes. The pipeline has been used for several genome annotations including several basidiomycetes [19, 21-35]. This demonstrated that significant differences exist in the plant polysaccharides degrading potential of *Schizophyllum commune*, *Coprinopsis cinerea*, *Laccaria bicolor*, *Postia placenta*, *Phanerochaete chrysosporium*, *Cryptococcus neoformans* and *Ustilago maydis* [25]. Overall, the genome of *C. neoformans* contained the lowest number of genes encoding putative plant cell wall (PCW) degrading enzymes, followed by the genomes of *U. maydis* and *L. bicolor*, which can be linked to their lifestyle as a animal/human pathogen, a biotrophic plant pathogen and a mycorrhizae, respectively. The saprobic basidiomycetes all contain significantly higher numbers of genes encoding putative PCW enzymes.

Table 1. Fungal enzymatic activities involved in plant polysaccharide degradation. The enzymes are sorted alphabetically on the enzyme code. Modified from [19].

Enzyme class	Enzyme code	Substrate	CAZy families
α -L-arabinofuranosidase	ABF	Xyloglucan, xylan, pectin	GH51,54
endoarabinanase	ABN	Pectin	GH43
exoarabinanase	ABX	Pectin	GH93
α -L-fucosidase	AFC	Xyloglucan	GH29,95
α -1,4-D-glucosidase	AGD	Starch	GH31
α -1,4-D-galactosidase	AGL	Xyloglucan, xylan, galactomannan	GH27,36
α -glucuronidase	AGU	Xylan	GH67,115
α -amylase	AMY	Starch	GH13
acetyl xylan esterase	AXE	Xylan	CE1
arabinoxylan arabinofuranohydrolase	AXH	Xylan	GH62
α -D-xylosidase	AXL	Xyloglucan	GH31
β -1,4-D-glucosidase	BGL	Cellulose	GH1,3
β -1,4-D-xylosidase	BXL	Xylan, pectin	GH3,39,43
cellobiohydrolase	CBH	Cellulose	GH6,7
β -1,4-D-endoglucanase	EGL	Cellulose	GH5,7,12,61
feruloyl esterase	FAE	Xylan, pectin	CE1
β -1,4-endogalactanase	GAL	Pectin	GH53
Glucuronyl esterase	GE	Xylan	CE15
Glucoamylase	GLA	Starch	GH15
β -1,6-endogalactanase	GLN	Pectin	
galactomannan acetyl esterase	GMAE	Galactomannan	
Endo-inulinase	INU	Inulin	GH32
Exo-inulinase	INX	Inulin	GH32
β -1,4-D-galactosidase	LAC	Xyloglucan, xylan, pectin, galactomannan	GH2,35
β -1,4-D-endomannanase	MAN	Galactomannan	GH5,26
β -1,4-D-mannosidase	MND	Galactomannan	GH2
pectin acetyl esterase	PAE	Pectin	
pectin lyase	PEL	Pectin	PL1
rhamnogalacturonan lyase	RGL	Pectin	PL4,11
endopolygalacturonase	PGA	Pectin	GH28
exopolygalacturonase	PGX	Pectin	GH28
pectate lyase	PLY	Pectin	PL1,3,9
pectin methyl esterase	PME	Pectin	CE8
rhamnogalacturonan acetyl esterase	RGAE	Pectin	CE12
rhamnogalacturonan galaturonohydrolase	RGX	Pectin	GH28
/ exorhamnogalacturonase			
α -rhamnosidase / rhamnogalacturonan rhamnohydrolase	RHA	Pectin	GH78
rhamnogalacturonan hydrolase / endorhamnogalacturonase	RHG	Pectin	GH28
Invertase / fructofuranosidase	SUC	Inulin	GH32
d-4,5 unsaturated -glucuronyl hydrolase	UGH	Pectin	GH88
unsaturated rhamnogalacturonan hydrolase	URH	Pectin	GH105
β -1,4-exogalactanase	XFG	Pectin	
xyloglucan-active β -1,4-D-endoglucanase	XEG	Xyloglucan	GH12,74
xyloglucan acetylesterase	XGAE	Xyloglucan	
β -1,4-D-endoxylanase	XLN	Xylan	GH10,11
β -1,6-exogalactanase	XSG	Pectin	
β -1,3-exogalactanase	XTG	Pectin	

EXPANSION OF BASIDIOMYCETE GENOMES

Basidiomycete species form a relative small section among the fungi for which genome annotations have been published as could be seen from the comparison mentioned above. This prevents in depth comparisons as have been published for ascomycete species as the coverage of the basidiomycete tree of line is insufficient to draw conclusions on evolutionary changes in these species related to plant polysaccharide degradation. However, a much larger set of basidiomycete genomes is currently in progress which will soon enable such detailed studies (Table 2). With the rapidly reducing costs of genome sequencing, this number of genomes is likely to grown exponentially over the next years.

Table 2. Basidiomycete genomes that are finished or in progress.

Species	Order	Sequencing Institute
<i>Agaricus bisporus</i> var <i>bisporus</i>	Agaricomycotina	JGI
<i>Agaricus bisporus</i> var. <i>burnettii</i>	Agaricomycotina	JGI
<i>Auricularia delicata</i>	Agaricomycotina	JGI
<i>Ceriporiopsis subvermispora</i>	Agaricomycotina	JGI
<i>Coniophora puteana</i>	Agaricomycotina	JGI
<i>Coprinopsis cinerea</i>	Agaricomycotina	Broad
<i>Cryptococcus neoformans</i>	Agaricomycotina	Broad
<i>Dacryopinax</i> sp.	Agaricomycotina	JGI
<i>Dichomitus squalens</i>	Agaricomycotina	JGI
<i>Fomitiporia mediterranea</i>	Agaricomycotina	JGI
<i>Fomitopsis pinicola</i>	Agaricomycotina	JGI
<i>Ganoderma</i> sp.	Agaricomycotina	JGI
<i>Gloeophyllum trabeum</i>	Agaricomycotina	JGI
<i>Hemileia vastatrix</i>	Pucciniomycotina	Genoscope
<i>Heterobasidion annosum</i>	Agaricomycotina	JGI
<i>Laccaria bicolor</i>	Agaricomycotina	JGI
<i>Malassezia globosa</i>	Ustilaginomycotina	JGI
<i>Melampsora laricis-populina</i>	Pucciniomycotina	JGI
<i>Microbotryum violaceum</i>	Pucciniomycotina	Genoscope
<i>Phanerochaete carnosa</i>	Agaricomycotina	JGI
<i>Phanerochaete chrysosporium</i>	Agaricomycotina	JGI
<i>Phlebia brevispora</i>	Agaricomycotina	JGI
<i>Pleurotus ostreatus</i>	Agaricomycotina	JGI
<i>Postia placenta</i>	Agaricomycotina	JGI
<i>Puccinia graminis</i>	Pucciniomycotina	Broad
<i>Punctularia strigosozonata</i>	Agaricomycotina	JGI
<i>Rhodotorula graminis</i>	Pucciniomycotina	JGI
<i>Schizophyllum commune</i>	Agaricomycotina	JGI
<i>Serpula lacrymans</i>	Agaricomycotina	JGI
<i>Sporobolomyces roseus</i>	Pucciniomycotina	JGI
<i>Stereum hirsutum</i>	Agaricomycotina	JGI
<i>Trametes versicolor</i>	Agaricomycotina	JGI
<i>Tremella mesenterica</i> Fries	Agaricomycotina	JGI
<i>Ustilago maydis</i>	Ustilaginomycotina	Broad
<i>Wolfiporia cocos</i>	Agaricomycotina	JGI

GROWTH PROFILING OF FUNGI VALIDATES DIFFERENCES IN FUNGAL GENOME SEQUENCES RELATED TO PLANT POLYSACCHARIDE DEGRADATION

Recently a novel fungal database has been initiated (www.fung-growth.org) to host growth profiles on a set of 35 carbon sources related to plant biomass. These growth profiles have already been used to provide biological support for differences observed in ascomycete genomes with respect to plant polysaccharide degradation [19, 22-24, 27]. The database currently contains growth profiles for more than 100 fungal species with genome sequences that are finished or in progress and has already shown significant differences in the ability of basidiomycete fungi to use various plant polysaccharides. These comparisons are part of several basidiomycete genome papers that are currently submitted or in progress and will provide a better understanding of the evolutionary changes in basidiomycete fungi.

CONCLUDING REMARKS

Considering the importance of plant biomass as a carbon source for many fungi, a better understanding of the diversity of fungi with respect to carbon utilisation will provide a significant increase in our understanding of fungal evolution. The expansion of basidiomycete genome sequences and other datasets (e.g. transcriptomics, proteomics) will enable studies into these fungi to much higher level and is likely to result in new strategies for improving mushroom cultivation.

REFERENCES

- [1] de Vries, R. P., and J. Visser. 2001. *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides. *Microb. Mol. Biol. Rev.* 65:497-522.
- [2] Kolpak, F. J., and J. Blackwell. 1976. Determination of the structure of cellulose II. *Macromolecules* 9:273-278.
- [3] van den Brink, J., and R. P. de Vries. 2011. Fungal enzyme sets for plant polysaccharide degradation. *Appl Microbiol Biotechnol.*
- [4] Ebringerová, A., and T. Heinze. 2000. Xylan and xylan derivatives - biopolymers with valuable properties, 1 Naturally occurring xylans structures, isolation procedures and properties. *Macromol. Rapid Commun.* 21:542-556.
- [5] Wilkie, K. C. B. 1979. The hemicelluloses of grasses and cereals. *Adv. Carbohydr. Chem. Biochem.* 36:215-264.
- [6] Brillouet, J.-M., and J. P. Joseleau. 1987. Investigation of the structure of a heteroxylan from the outer pericarp (beeswing bran) of wheat kernel. *Carbohydr. Res.* 159:109-126.
- [7] Huisman, M. M. H., et al. 2000. Glucuronoarabinoxylans from maize kernel cell walls are more complex than those from sorghum kernel cell walls. *Carbohydr. Pol.* 43:269-279.
- [8] Schooneveld-Bergmans, M. E. F., et al. 1998. Extraction and partial characterization of feruloylated glucuronoarabinoxylans from wheat bran. *Carbohydr. Polymers* 35:39-47.
- [9] Saulnier, L., et al. 1995. Isolation and partial characterization of feruloylated oligosaccharides from maize bran. *Carbohydr. Res.* 272:241-253.
- [10] Smith, M. M., and R. D. Hartley. 1983. Occurrence and nature of ferulic acid substitution of cell wall polysaccharides in graminaceous plants. *Carbohydr. Res.* 118:65-80.

- [11] Wende, G., and S. C. Fry. 1997. O-feruloylated, O-acetylated oligosaccharides as side-chains of grass xylans. *Phytochem.* 44:1011-1018.
- [12] Vincken, J.-P., et al. 1997. Two general branching patterns of xyloglucan, XXXG and XXGG. *Plant Physiol.* 114:9-13.
- [13] Dey, P. M. 1978. Biochemistry of plant galactomannans. *Adv. Carbohydr. Chem. Biochem.* 35:3411-376.
- [14] de Vries, J. A., et al. 1982. Enzymic degradation of apple pectins. *Carbohydr. Polymers* 2:25-33.
- [15] Perez, S., et al. 2000. The three-dimensional structures of the pectic polysaccharides. *Plant Physiol. Biochem.* 38:37-55.
- [16] Hoover, R., and F. W. Sosulski. 1991. Composition, structure, functionality, and chemical modification of legume starches: a review. *Can J Physiol Pharmacol* 69:79-92.
- [17] Ritsema, T., and S. Smeeckens. 2003. Fructans: beneficial for plants and humans. *Curr Opin Plant Biol* 6:223-30.
- [18] Verbeke, D., et al. 2003. Exudate gums: occurrence, production, and applications. *Appl Microbiol Biotechnol* 63:10-21.
- [19] Coutinho, P. M., et al. 2009. Post-genomic insights into the plant polysaccharide degradation potential of *Aspergillus nidulans* and comparison to *Aspergillus niger* and *Aspergillus oryzae*. *Fungal Genet Biol* 46 Suppl 1:S161-S169.
- [20] Cantarel, B. L., et al. 2009. The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res* 37:D233-8.
- [21] Andersen, M. R., et al. Comparative genomics of citric-acid-producing *Aspergillus niger* ATCC 1015 versus enzyme-producing CBS 513.88. *Genome Res* 2011.
- [22] Espagne, E., et al. 2008. The genome sequence of the model ascomycete fungus *Podospora anserina*. *Genome Biol.* 9:R77 (1-22).
- [23] Goodwin, S. B., et al. 2011. Finished Genome of the Fungal Wheat Pathogen *Mycosphaerella graminicola* Reveals Dispensome Structure, Chromosome Plasticity, and Stealth Pathogenesis. *PLoS Genet* 7:e1002070.
- [24] Levesque, C. A., et al. 2010. Genome sequence of the necrotrophic plant pathogen *Pythium ultimum* reveals original pathogenicity mechanisms and effector repertoire. *Genome Biol* 11:R73.
- [25] Ohm, R. A., et al. 2010. Genome sequence of the model mushroom *Schizophyllum commune*. *Nat Biotechnol.*
- [26] Pel, H. J., et al. 2007. Genome sequencing and analysis of the versatile cell factory *Aspergillus niger* CBS 513.88. *Nat Biotechnol* 25:221-31.
- [27] Battaglia, E., et al. 2011. Carbohydrate-active enzymes from the zygomycete fungus *Rhizopus oryzae*: a highly specialized approach to carbohydrate degradation depicted at genome level. *BMC Genomics* 12:38.
- [28] Coleman, J. J., et al. 2009. The genome of *Nectria haematococca*: contribution of supernumerary chromosomes to gene expansion. *PLoS Genet* 5:e1000618.
- [29] Duplessis, S., et al. From the Cover: Obligate biotrophy features unraveled by the genomic analysis of rust fungi. *Proc Natl Acad Sci U S A* 108:9166-71.
- [30] Kubicek, C. P., et al. Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*. *Genome Biol* 12:R40.
- [31] Ma, L. J., et al. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464:367-73.

- [32] Martin, F., et al. 2008. The genome of *Laccaria bicolor* provides insights into mycorrhizal symbiosis. *Nature* 452:88-92.
- [33] Martinez, D., et al. 2008. Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocrea jecorina*). *Nat Biotechnol* 26:553-60.
- [34] Martinez, D., et al. 2009. Genome, transcriptome, and secretome analysis of wood decay fungus *Postia placenta* supports unique mechanisms of lignocellulose conversion. *Proc Natl Acad Sci U S A* 106:1954-9.
- [35] Martinez, D., et al. 2004. Genome sequence of the lignocellulose degrading fungus *Phanerochaete chrysosporium* strain RP78. *Nat. Biotechnol.* 22:695-700.